


OpenCLIM Local Climate Risk Reports

Local insights from national analysis



What's included?

Agriculture

- Oil seed rape potential yield (units = tonnes per hectare, t/ha)
- Grass potential yield (t/ha)
- Wheat potential yield (t/ha)

These metrics indicate areas where a crop is likely to increase/decrease in yield due to climate-forced changes in temperature and water availability. 1 km grids.

Biodiversity

- Conservation potential (relative units)
- Restoration potential (relative units)
- Urban green space potential (relative units)

These metrics indicate the relative biodiversity impact in a location based on the richness of species remaining. 100 m grids (resampled from 10 m original analysis).

Heat Stress

- Heat-related mortality (units = mean deaths, cumulative deaths)


This metric projects the number of excess deaths due to prolonged or extreme warm weather conditions. 12 km grid.

Hydrology

- Drought duration (units = cumulative months per 30 year period)
- 1-in-10-year return period flow (units = flow rate in metres cubed per second, m³/s; difference, %)
- 1-in-100-year return period flow (units = flow rate in metres cubed per second, m³/s; difference, %)

These river flow-rate metrics are shown as a proxy for flooding and should be used alongside e.g. Environment Agency flood indicators. They are gridded at 1km for whole catchments.

Greater London | Potential Yield | Oil Seed Rape



Key Points

Potential yield (tonnes per hectare) shows change in potential oil seed rape yield at 1km resolution, due to heat limitation and water limitation under baseline, 2°C, and 4°C warming scenarios.

Scenarios shown include the CO2 fertilisation effect (enhanced plant productivity).


A modest increase in potential yield is projected at 2°C and 4°C for most of the Greater London area.

Local Summary

Minimum, mean and maximum potential yield (t/ha) for the Greater London region at baseline, 2°C and 4°C warming scenarios.

County	Mean			Minimum			Maximum		
	Baseline	2°C	4°C	Baseline	2°C	4°C	Baseline	2°C	4°C
City of London									
Greater London	8.4	9.8	11	7.5	7.8	7.6	8.8	10.5	10

Greater London | Potential Yield - difference | Oil Seed Rape



Key Points

Potential yield (tonnes per hectare) shows change in potential oil seed rape yield at 1km resolution, due to heat limitation and water limitation under baseline scenarios with the difference from baseline at 2°C and 4°C warming level scenarios.

Scenarios shown include the CO₂ fertilisation effect (enhanced plant productivity).


A modest increase in potential yield is projected at 2°C and 4°C for most of the Greater London area.

Local Summary

Minimum, mean and maximum potential yield (t/ha) for the Greater London region at baseline, 2°C and 4°C warming scenarios, with the difference from the baseline mean for 2°C and 4°C warming scenarios.

County	Mean			Minimum			Maximum			Difference	
	Baseline	2°C	4°C	Baseline	2°C	4°C	Baseline	2°C	4°C	2°C	4°C
City of London											
Greater London	8.4	9.8	11	7.5	7.8	7.6	8.8	10.5	10	1.4	2.6

Greater London | Potential Yield | Grass



Key Points

Potential yield (tonnes per hectare) shows change in potential grass yield at 1km resolution, due to heat limitation and water limitation under baseline, 2°C, and 4°C warming scenarios.

Scenarios shown include the CO2 fertilisation effect (enhanced plant productivity).


A modest increase in potential yield is projected at 2°C and 4°C for most of the Greater London area.

Local Summary

Minimum, mean and maximum potential yield (t/ha) for the Greater London region at baseline, 2°C and 4°C warming scenarios.

County	Mean			Minimum			Maximum		
	Baseline	2°C	4°C	Baseline	2°C	4°C	Baseline	2°C	4°C
City of London									
Greater London	6.6	7.4	8.9	6.3	7	8.4	6.8	7.7	9.2

Greater London | Potential Yield - difference | Grass



Key Points

Potential yield (tonnes per hectare) shows change in potential grass yield at 1km resolution, due to heat limitation and water limitation under baseline scenarios with the difference from baseline at 2°C and 4°C warming level scenarios.

Scenarios shown include the CO₂ fertilisation effect (enhanced plant productivity).


A modest increase in potential yield is projected at 2°C and 4°C for most of the Greater London area.

Local Summary

Minimum, mean and maximum potential yield (t/ha) for the Greater London region at baseline, 2°C and 4°C warming scenarios, with the difference from the baseline mean for 2°C and 4°C warming scenarios.

County	Mean			Minimum			Maximum			Difference	
	Baseline	2°C	4°C	Baseline	2°C	4°C	Baseline	2°C	4°C	2°C	4°C
City of London											
Greater London	6.6	7.4	8.9	6.3	7	8.4	6.8	7.7	9.2	0.8	2.3

Greater London | Potential Yield | Wheat



Key Points

Potential yield (tonnes per hectare) shows change in potential wheat yield at 1km resolution, due to heat limitation and water limitation under baseline, 2°C, and 4°C warming scenarios.

Scenarios shown include the CO2 fertilisation effect (enhanced plant productivity).


There is a slight increase in potential yield projected at 2°C and 4°C for most of the Greater London area.

Local Summary

Minimum, mean and maximum potential yield (t/ha) for the Greater London region at baseline, 2°C and 4°C warming scenarios.

County	Mean			Minimum			Maximum		
	Baseline	2°C	4°C	Baseline	2°C	4°C	Baseline	2°C	4°C
City of London									
Greater London	14	14.4	13.6	8.1	7.5	6.6	14.6	17	17.8

Greater London | Potential Yield - difference | Wheat



Key Points

Potential yield (tonnes per hectare) shows change in potential wheat yield at 1km resolution, due to heat limitation and water limitation under baseline scenarios with the difference from baseline at 2°C and 4°C warming level scenarios.

Scenarios shown include the CO2 fertilisation effect (enhanced plant productivity).


There is slight increase in potential yield projected at 2°C and 4°C for most of the Greater London area.

Local Summary

Minimum, mean and maximum potential yield (t/ha) for the Greater London region at baseline, 2°C and 4°C warming scenarios, with the difference from the baseline mean for 2°C and 4°C warming scenarios.

County	Mean			Minimum			Maximum			Difference	
	Baseline	2°C	4°C	Baseline	2°C	4°C	Baseline	2°C	4°C	2°C	4°C
City of London											
Greater London	14	14.4	13.6	8.1	7.5	6.6	14.6	17	17.8	0.4	-0.4

Greater London | Conservation | Warming Levels



Key Points

A relative scoring is shown for an area's suitability for Conservation, based on a metric of species richness remaining.

Under 2°C and 4°C warming scenarios, a decline in suitability is projected, with mean values shown in the Table (right).


Under 4°C of warming, only one area is projected to retain a maximum suitability of 100 (range = 60 to 100), while the mean rarely exceeds 50, showing the importance of limiting warming to 2°C or less.

Local Summary

Minimum, mean and maximum conservation potential (%) for Greater London at baseline, 2°C and 4°C warming scenarios.

County	Mean			Minimum			Maximum		
	Baseline	2°C	4°C	Baseline	2°C	4°C	Baseline	2°C	4°C
City of London	100	74		100	74		100	74	
Greater London	26.7	20.3	16.4	0	0	0	100	88	68

Greater London | Restoration | Warming Levels



Key Points

A relative scoring is shown for an area's suitability for Restoration, based on a metric of species richness remaining.

Under 2°C and 4°C warming scenarios, a decline in suitability is projected, with mean values shown in the Table (right).


Under 4°C of warming no areas remain with a projected maximum suitability of 100 (range = 60 to 95), showing the importance of limiting warming to 2°C or less.

Local Summary

Minimum, mean and maximum restoration potential (%) for Greater London at baseline, 2°C and 4°C warming scenarios.

County	Mean			Minimum			Maximum		
	Baseline	2°C	4°C	Baseline	2°C	4°C	Baseline	2°C	4°C
City of London	0	0		0	0		0	0	
Greater London	73.3	55.1	43.9	0	0	0	100	88	68

Greater London | Urban Green | Warming Levels



Key Points

A relative scoring is shown for an area's suitability for Urban Green Space, based on a metric of species richness remaining.

Under 2°C and 4°C warming scenarios, a decline in suitability is projected, with mean values shown in the Table (right).

Under 4°C of warming there are no areas remaining with a maximum suitability of 100 (range = 60 to 95), showing the importance of limiting warming to 2°C or less.

Local Summary

Minimum, mean and maximum urban greenspace potential (%) for Greater London at baseline, 2°C and 4°C warming scenarios.

County	Mean			Minimum			Maximum		
	Baseline	2°C	4°C	Baseline	2°C	4°C	Baseline	2°C	4°C
City of London	99.7	74.2	60	0	0	60	100	75	60
Greater London	70.9	53.4	42.7	0	0	0	100	88	68

Greater London | Heat Mortality | Combined Future Scenarios

Number of Deaths

a) Baseline




0
1-2
2-5
5-10
10-25

b) SSP2, 2°C, 2050

0
2-5
5-10
10-25
25-50
50-100

c) SSP2, 4°C, 2080

2-5
5-10
10-25
25-50
50-100
100-200
>200



Key Points

Heat mortality shows average deaths per year at 12km resolution, under future scenarios combining warming (2°C, 4°C), socioeconomics (SSP2, SSP4), and population (2050, 2080).

An increase in heat mortality is projected under 2°C and 4°C scenarios under SSP2. Additional population in 2050 and 2080 also increase mortality.

The climate model ensemble shows a range of outcomes, summarised by the 10th to 90th percentile range (bottom Table, right).

Local Summary

Mean deaths per year and cumulative deaths in Greater London for baseline and future scenarios.

County	Baseline		SSP2 2050 2°C			SSP2 2080 4°C		
	Mean	Cumulative	Mean	Cumulative	Difference	Mean	Cumulative	Difference
Greater London	4.8	110.4	27.3	628	22.5	89.8	2066.2	85

Mean deaths (death/yr) for each future scenario and the climate model ensemble range between 10th and 90th percentile.

County	Baseline Mean	10th - 90th percentile	SSP2 2050 2°C Mean	10th - 90th percentile	SSP2 2080 4°C Mean	10th - 90th percentile
Greater London	4.8	0.6 - 18.1	27.3	4.5 - 102.4	89.8	14.1 - 325.5

Greater London | Heat Mortality | Combined Future Scenarios

Number of Deaths

a) Baseline




0
1-2
2-5
5-10
10-25

b) SSP4, 2050, 2°C

0
2-5
5-10
10-25
25-50
50-100

c) SSP4, 2080, 4°C

1-2
5-10
10-25
25-50
50-100
100-200
>200



Key Points

Heat mortality shows average deaths per year at 12km resolution, under future scenarios combining warming (2°C, 4°C), socioeconomics (SSP2, SSP4), and population (2050, 2080).

An increase in heat mortality is projected under 2°C and 4°C scenarios under SSP4. Additional population in 2050 and 2080 also increase mortality.

The climate model ensemble shows a range of outcomes, summarised by the 10th to 90th percentile range (bottom Table, right).

Local Summary


Mean deaths per year and cumulative deaths in Greater London for baseline and future scenarios.

County	Baseline		SSP4 2050 2°C			SSP4 2080 4°C		
	Mean	Cumulative	Mean	Cumulative	Difference	Mean	Cumulative	Difference
Greater London	4.8	110.4	24.8	569.9	20	71.6	1646	66.8

Mean deaths (death/yr) for each future scenario and the climate model ensemble range between 10th and 90th percentile.

County	Baseline Mean	10th - 90th percentile	SSP4 2050 2°C Mean	10th - 90th percentile	SSP4 2080 4°C Mean	10th - 90th percentile
Greater London	23.5	4.2 - 84.6	24.8	4.2 - 91.7	71.6	11.9 - 253.6

Greater London | Drought Duration | Warming Levels



Key Points

Drought duration is a low-flow metric representing the average cumulative duration of drought projected within a future 30-year period.

Gridded 1km results are modelled at catchment scale. Coastal or tidally influenced catchments are not modelled.

Nationally for 2°C and 4°C warming scenarios, most catchments are projected to experience an increase in drought duration, particularly in southern and eastern areas.

The climate model ensemble shows a range of possible future outcomes, summarised by the 10th to 90th percentile range (bottom table).

Local Summary


Median, minimum, and maximum drought duration (months/30-yr) for baseline scenario in Greater London, and the percentage change from baseline for a 2°C and 4°C warming scenario.

County	Median			Minimum			Maximum		
	Baseline	2°C	4°C	Baseline	2°C	4°C	Baseline	2°C	4°C
City of London									
Greater London	2.7	3.1	3.6	2.2	2.4	2.7	3.6	7.1	13.3

Median drought duration (months/30-yr) for baseline scenario in Greater London, the climate model ensemble range between 10th and 90th percentile, and the percentage change from baseline for a 2°C and 4°C warming scenario.

County	Baseline	10th - 90th	2°C	10th - 90th	4°C	10th - 90th
	Median	percentile	Median	percentile	Median	percentile
City of London		-		-		-
Greater London	2.7	1.8 - 5.3	3.1	2.1 - 12.4	3.6	2.4 - 24.2

Greater London | 10 year Return Period Flow | Warming Levels



Key Points

The 1-in-10-year return period flow is a high-flow rate metric with a 10% annual probability of occurring. It is a proxy for a low probability, low magnitude flood event.

Nationally for 2°C of warming, most catchments are projected to experience 5% to 30% increase in flows, with little additional increase for most catchments at 4°C. Flow is projected to decrease for some central & eastern catchments.

The climate model ensemble shows a range of outcomes which is summarised by the 10th to 90th percentile range (bottom table).

Local Summary

Median, minimum, and maximum flow rate (m³/s) for baseline scenario in Greater London, and the percentage change from baseline for a 2°C and 4°C warming scenario.


County	Median			% change			Minimum			Maximum		
	Baseline	2°C	4°C	Baseline	2°C	4°C	Baseline	2°C	4°C	Baseline	2°C	4°C
City of London												
Greater London	5.5	9.6	22.8	2.7	-15.4	-20.1	424.2	22.2	45.9			

Median flow rate (m³/s) for baseline scenario in Greater London, the climate model ensemble range between 10th and 90th percentile, and the percentage change from baseline for a 2°C and 4°C warming scenario.


County	Baseline Median	10th - 90th percentile	2°C % change	4°C % change
City of London		-		
Greater London	5.5	2.2 - 483.8	9.6	22.8

Greater London | 100 year Return Period Flow | Warming Levels



Flow Rate

a) Baseline (m³/s)

b) 2°C % change



c) 4°C % change



Key Points

The 1-in-100-year return period flow is a high-flow rate metric with a 1% annual probability of occurring. It is a proxy for a low probability, high magnitude flood event.

Nationally for 2°C warming, most catchments are projected to experience 5% to 30% increase in flows, while at 4°C more catchments are projected to increase flow >60%. Flow is projected to decrease for some catchments in the East.

The climate model ensemble shows a range of outcomes which is summarised by the 10th to 90th percentile range (bottom table).

Local Summary

Median, minimum, and maximum flow rate (m³/s) for baseline scenario in Greater London, and the percentage change from baseline for a 2°C and 4°C warming scenario.

County	Median			% change			Minimum			Maximum		
	Baseline	2°C	4°C	Baseline	2°C	4°C	Baseline	2°C	4°C	Baseline	2°C	4°C
City of London												
Greater London	7.7	18.6	47.2	3	-19	-16.3	1038.4	50.9	91.3			

Median flow rate (m³/s) for baseline scenario in Greater London, the climate model ensemble range between 10th and 90th percentile, and the percentage change from baseline for a 2°C and 4°C warming scenario.

County	Baseline Median	10th - 90th percentile	2°C % change	4°C % change
City of London		-		
Greater London	7.7	2.5 - 856.3	18.6	47.2

Further information

OpenCLIM disclaimer / intended use statement

Exclusion of Liability

Whilst every effort has been made to ensure the accuracy of information presented, the OpenCLIM team and partner institutions disclaim all responsibility for and accept no liability for any errors or losses caused by any inaccuracies in such information or the consequences of any person acting or refraining from acting or otherwise relying on such information.

Your use of information provided by OpenCLIM is at your own risk. Please read any warnings given about the limitations of the information.

The OpenCLIM team and partner institutions give no warranty as to the quality or accuracy of the information or its suitability for any use. All implied conditions relating to the quality or suitability of the information, and all liabilities arising from the supply of the information (including any liability arising in negligence) are excluded to the fullest extent permitted by law.

The OpenCLIM team and partner institutions give no warranty as to the accuracy or completeness of data or images in the form in which they are cached or downloaded to your computer, as they may be affected by online conditions beyond our control.

Intended-Use

The OpenCLIM framework is a national-scale set of workflows and datasets for describing climate change risks/opportunities and for assessing adaptation needs. Data are at resolutions appropriate for investigating climate risks at national and sub-national scales. OpenCLIM is not intended to provide information about climate risks at the scale of individual properties or streets and should not be used for such micro analysis.

Limitations

- Data, model results and other information or related records have been produced for a specific purpose, and that may affect the type and completeness of the data and interpretation. The nature and purpose of data collection and model simulation may render it unsuitable for certain applications/uses. You must verify the suitability of the material for your intended usage.
- The data, information and related records within the OpenCLIM framework should not be taken as a substitute for specialist interpretations, professional advice and/or detailed site investigations. You must seek professional advice before making technical interpretations on the basis of the materials provided.

Sharing and re-use

OpenCLIM results are shared under a creative commons licence ([CCBY 4.0](https://creativecommons.org/licenses/by/4.0/)) and may be re-used with appropriate credit to the original creators.



Front cover image credits: Don Lodge and Bruno Martins on [Unsplash.com](https://unsplash.com/).

Maps contain public sector information (administrative boundaries) licensed under the [Open Government Licence v3.0](https://www.gov.uk/government/publications/open-government-licence-v3-0).

